



## **Risks of non-cancer diseases at low and intermediate dose levels in the French cohort of nuclear workers: updated follow-up 1968-2014**

Klervi Leuraud, Dominique Laurier, Lucie Fournier, Sylvaine Caer-Lorho, Eric Samson

### **► To cite this version:**

Klervi Leuraud, Dominique Laurier, Lucie Fournier, Sylvaine Caer-Lorho, Eric Samson. Risks of non-cancer diseases at low and intermediate dose levels in the French cohort of nuclear workers: updated follow-up 1968-2014. 6th European Radiological Protection Research Week (ERPW 2022), Oct 2022, Estoril, Portugal. irsn-03940244

**HAL Id: irsn-03940244**

**<https://irsnn.hal.science/irsnn-03940244>**

Submitted on 16 Jan 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright





# Risks of non-cancer diseases at low and intermediate dose levels in the French cohort of nuclear workers: updated follow-up 1968-2014

Klervi LEURAUD<sup>1</sup>, Dominique LAURIER<sup>1</sup>, Lucie FOURNIER<sup>1</sup>, Sylvaine CAËR-LORHO<sup>1</sup>, Eric SAMSON<sup>1</sup>, Olivier LAURENT<sup>1</sup>  
1. Institute for Radiological Protection and Nuclear Safety (IRSN), Fontenay-aux-Roses, France

## INTRODUCTION

### Context

- Radiation-associated risks of non-cancer diseases remain poorly understood in the low dose/low dose-rate range.
- Populations of nuclear workers are particularly relevant to study health effects of protracted exposures to low doses at low dose-rates of ionizing radiation (IR).

### Objective

- To report on risk of death from non-cancer diseases of nuclear workers badge-monitored for external IR exposure and followed-up through several decades
- To strengthen the basis for radiation protection.

## MATERIALS AND METHODS

- The SELTINE cohort of nuclear workers employed by CEA, Orano, or EDF companies in France was followed-up for mortality from 1968 to 2014.
- Cumulative doses due to occupational external IR (mostly photons) were reconstructed individually and expressed either in  $H_p(10)$  dose equivalents in Sievert (Sv) or absorbed doses to organs of interest in Gray (Gy).
- Estimates of excess relative rate (ERR) per Sv or Gy have been calculated using Poisson regression methods and a linear ERR model:
  - ERR were estimated only for males who represent 87% of the cohort;
  - the background mortality rate was modeled as a function of age, calendar year, company, duration of employment, and socioeconomic status;
  - cumulative doses were lagged by 10 years to allow for a latent period in radiation effect.

## RESULTS

Characteristics of the SELTINE cohort of nuclear workers in France (males only)

Number of male workers	69,487	
Number of accrued person-years	2,202,492	
Follow-up (1968-2014), in years (SD)		
Mean duration	31.7	(11.4)
Mean age at end of follow-up	62.8	(13.6)
Vital status on 12/31/2014, n (%)		
Alive	55,134	(79.3)
Deceased	14,131	(20.4)
Lost to follow-up	222	(0.3)
Employment		
Mean age at hiring, in years (SD)	26.4	6.3
Mean duration, in years (SD)	26.5	(9.0)
Socioeconomic status, n (%)		
Managers and engineers	13,466	(19.4)
Administrative employees	2,909	(4.2)
Skilled workers	40,537	(58.3)
Unskilled workers	11,787	(17.0)
Unknown	788	(1.1)
Cumulative Hp(10) dose, in mSv		
Number of exposed* workers (%)	49,995	(71.9)
Mean (SD), whole cohort	17.7	(39.7)
Mean (SD), exposed* workers only	24.7	(44.9)

SD: standard deviation; \* with at least one positive recorded dose

Excess relative rate (ERR) of death from non-cancer diseases per unit of 10-year lagged cumulative dose in males only, SELTINE cohort, 1968-2014

Cause of death	Target tissue	Observed deaths	ERR/Gy	95% CI	p-value
Non-cancer diseases	H <sub>p</sub> (10)	8,577	0.02	-0.49; 0.57	0.952
Diabetes mellitus	H <sub>p</sub> (10)	169	-0.91	ne; 2.62	0.536
Mental and behavioral disorders	Brain	235	2.74	-1.95; 10.33	0.316
Diseases of the nervous system	Brain	531	3.47	-0.48; 8.56	0.091
Dementia, Alzheimer’s disease	Brain	269	9.62	3.05; 18.68	<b>0.002</b>
Parkinson’s disease	Brain	124	-1.30	ne; 7.44	0.711
Circulatory diseases	H <sub>p</sub> (10)	3261	0.09	-0.72; 1.03	0.834
Ischemic diseases	H <sub>p</sub> (10)	1,258	-0.23	ne; 1.38	0.763
Cerebrovascular diseases	H <sub>p</sub> (10)	684	1.41	ne; 4.05	0.194
Hypertensives diseases	H <sub>p</sub> (10)	120	2.31	ne; 9.98	0.374
Respiratory diseases	Lung	558	0.22	ne; 3.71	0.884
Chronic obstructive pulmonary diseases	Lung	164	0.15	ne; 7.52	0.958
Digestive diseases	Colon	583	0.96	-1.72; 4.58	0.531
Cirrhosis	Liver	166	3.72	ne; 13.64	0.288

\* estimated from a linear model, adjusted on calendar year, age, sex, company, duration of employment, and socioeconomic status; CI: likelihood-based confidence interval; NC: convergence not achieved; *ne*: not estimated.

### Summary of results

- No association was observed for most of outcomes
- Only mortality from dementia and Alzheimer’s disease showed a statistically significant dose-risk association

## CONCLUSION

- The long follow-up makes this cohort informative with respect to the long-term risks following protracted exposure to low-dose ionizing radiation.
- The observed association between dementia mortality risk and ionizing radiation was not observed in previous analyses and should be viewed with caution
- The association between dementia mortality risk and IR should be further investigated by other studies

**Acknowledgments.** The construction of the French cohort was realized by the IRSN, with partial funding from Orano and EDF. This funding had no impact on the study design, analyses, or interpretation of results. IRSN thanks all persons from CEA, Orano, and EDF who cooperated in the elaboration of the French cohort by providing access to sources of administrative and dosimetric data for this study