

Tracing airborne micro-particles released by Al-industry using actinides

Laurent Pourcelot, Olivier Masson, Beatrice Boulet

▶ To cite this version:

Laurent Pourcelot, Olivier Masson, Beatrice Boulet. Tracing airborne micro-particles released by Al-industry using actinides. 3rd International Conference on Radioanalytical and Nuclear Chemistry (RANC-3), American Nuclear Society, May 2023, Budapest, Hungary. irsn-04113866

HAL Id: irsn-04113866 https://irsn.hal.science/irsn-04113866v1

Submitted on 1 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Tracing airborne micro-particles released by Al-industry using actinides

Laurent Pourcelot^{1*}, Olivier Masson¹, Béatrice Boulet²

Keywords: actinides, aerosols, tree leaves, red muds

Industry remains an important source of environmental contamination by heavy metals and metalloids which may pose a serious threat to the ecosystems and the population. Especially the alumina production from bauxite ore gives rise to large amount of liquid and solid wastes since the production of 1 kg of alumina involves same weight of solid residue, the well-known red muds enriched in iron and heavy metals. Up to now, without any valorization of by-products, the red muds are stored in artificial ponds which are a potential source of pollutants for the surroundings due to the uplift of red dust under strong wind conditions. The study of plants allows filling the gap of knowledge concerning air quality over a wide potentially contaminated area. Thus, the current work aims to compare the activity concentration of natural radionuclides (²³⁸U, decay products and ²³²Th) in plants leaves (*quercus robur* and lettuces), in grains (wheat) and leaf vegetables with those of aerosols and the potential sources of atmospheric particles from soils and above all the red dust emitted from the dried ponds and from the piles of bauxite.

The activity concentration of natural radionuclides such as ²³⁸U, some decay products and ²³²Th was determined in plants samples taken according to the two dominant wind directions, at plots located 100 to 1,500 m from the basins. Furthermore, the atmospheric particles were taken in the same areas, using a high-volume aerosol sampler. The increase of ²³²Th and ²³⁸U activity concentration in a few trees leaves (a factor 9 and 4, respectively) and in some aerosols samples is accompanied by a decrease of ²³⁸U/²³²Th activity ratio of these matrices. Such low ratio suggests that the airborne particles emitted by bauxite piles and red mud basins - those latter's are also characterized by low ²³⁸U/²³²Th - first contaminate the atmosphere and then the leaves surface, after deposition. Available at distance greater than 1,000 m from the ponds, locally produced foodstuffs do not show any excess of radionuclides, suggesting low influence of airborne micro particles from the alumina production. Thus, the actinides measured in leaf vegetables rather derived from the soil micro-particles deposited onto leaves than from an anthropogenic source.

Figures

Insert figures here, if applicable.

figure caption

Table

source/note

Acknowledgments

If applicable.

References

List of References here, if applicable.

You will receive a confirmation email upon submission. **Please check your spam folder as well** in case it does not appear in your mailbox.

¹Institut de Radioprotection et de Sûreté Nucléaire IRSN, Saint-Paul-lez-Durance, France

²Institut de Radioprotection et de Sûreté Nucléaire IRSN, Orsay, Country

^{*}E-mail: laurent.pourcelot@irsn.fr