



Radioactivity in the Marine Environment: Cosmogenic and Anthropogenic Radionuclides

Claudia Benitez-Nelson, Ken Buesseler, Minhan Dai, Michio Aoyama, Núria Casacuberta, Sabine Charmasson, José Marcus Godoy, Andy Johnson, Vladimir Maderich, Pere Masqué, et al.

► To cite this version:

Claudia Benitez-Nelson, Ken Buesseler, Minhan Dai, Michio Aoyama, Núria Casacuberta, et al.. Radioactivity in the Marine Environment: Cosmogenic and Anthropogenic Radionuclides. Limnology and Oceanography e-Lectures, 2018, 8 (1), pp.114-169. 10.1002/loe2.10008 . irsn-04558936

HAL Id: irsn-04558936

<https://irsn.hal.science/irsn-04558936>

Submitted on 25 Apr 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

Copyright

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Radioactivity in the Marine Environment

Cosmogenic and Anthropogenic Radionuclides


Claudia R. Benitez-Nelson

Ken Buesseler, Minhan Dai,
Michio Aoyama, Núria Casacuberta,
Sabine Charmasson, José Marcus Godoy,
Andy Johnson, Vladimir Maderich, Pere Masqué,
Willard Moore, Paul J. Morris, John N. Smith

Radionuclides in the Marine Environment

Radionuclides, of both natural and anthropogenic origins, can be used as CLOCKS of key processes (age and/or rates) in the oceans, mainly because they:

- Are ubiquitous in all compartments of the oceans
- Have a large range of half-lives (from seconds to billions of years)



Tracers in the Sea
Broecker and Peng, 1982

Three main sources of radionuclides to the Marine Environment:

1. **U-Th series radionuclides** – of primordial origin, occur naturally on land and in ocean, and produce a series of “daughter” radionuclides via radioactive decay.

Examples: ^{238}U , ^{234}Th , ^{232}Th , ^{210}Pb , $^{222}, 224, 226, 228\text{Ra}$ and ^{222}Rn .

2. **Cosmogenic Radionuclides** – continuously being created by cosmogenic rays that interact with materials in the atmosphere and on Earth.

Examples: ^3H , ^{14}C , ^7Be


3. **Anthropogenic radionuclides** – continuously being produced by humans.

Examples: ^3H , ^{14}C , ^{90}Sr , ^{137}Cs , ^{129}I , $^{238}, 239, 240\text{Pu}$


Note: Some radionuclides have both cosmogenic and anthropogenic sources (e.g., ^3H and ^{14}C).

Three main sources of radionuclides to the Marine Environment:

2) Cosmogenic (^{7}Be , ^{14}C)



3) Anthropogenic Radionuclides (^{90}Sr , ^{137}Cs)



Cosmogenic Radionuclides

Table 2

Cosmogenic nuclide source functions in oceans, for nuclides of half-lives > 10 days, arranged in order of increasing half-lives

Nuclide	Half-life	Principal target element(s)		Global average surface injection rate ^a (atoms/cm ² per min)	Integrated in situ oceanic production rate ^b
		Atmosphere	In ocean water		
³² P	14.3 days	Ar	Cl, S, K	5.82×10^{-3}	7.6×10^{-4}
³³ P	25.3 days	Ar	Cl, S, K	6.93×10^{-3}	2.9×10^{-4}
³⁷ Ar	35.0 days	Ar	K, Ca	9.10×10^{-6}	8.1×10^{-6}
⁷ Be	53.3 days	N,O	O	1.27	6.0×10^{-3}
³⁵ S	87.4 days	Ar	Cl, Ca, K	2.84×10^{-2}	5.1×10^{-4}
²² Na	2.6 years	Ar	Na	3.75×10^{-3}	3.9×10^{-4}
³ H	12.3 years	N,O	O, ² H	1.39×10^{-1}	1.2×10^{-2}
³² Si	~ 150 years	Ar	S, Ca	9.60×10^{-3}	2.5×10^{-5}
³⁹ Ar	269 years	Ar	K, Ca	2.00×10^{-1}	1.2×10^{-5}
¹⁴ C	5730 years	N,O	O	1.20×10^2	9.0×10^{-3}
⁴¹ Ca ^c	1.0×10^5 years	—	Ca	—	2.4×10^{-5e}
⁸¹ Kr	2.1×10^5 years	Kr	Sr	2.30×10^{-5}	1.9×10^{-8e}
³⁶ Cl ^d	3.0×10^5 years	Ar	Cl	6.60×10^{-2}	1.06×10^{-1e}
²⁶ Al	7.2×10^5 years	Ar	S, K, Ca	8.40×10^{-3}	6.8×10^{-6}
¹⁰ Be	1.6×10^6 years	N,O	O	2.70	1.8×10^{-3}

^a Based on atmospheric production rate estimates of Lal and Peters (1967). The mean stratosphere–troposphere exchange time is taken to be 2 years. Tropospheric residence time of isotopes that can be scavenged by wet precipitation is taken to be 40 days. Atmospheric residence times for ¹⁴C, ³⁹Ar, and ⁸¹Kr, for mixing with the mixed layer of the ocean, are taken to be 10, 270, and 170 years, respectively.

^b Lal et al. (1988).

^c Flux to oceans from rivers should be included to take into account production in rocks and soil by ⁴⁰Ca(*n*,γ) ⁴¹Ca reaction; this estimate is not given here because of large uncertainties in these calculations.

^d As above, due to ³⁵Cl(*n*,γ) ³⁶Cl reaction.


^e The in situ production of ⁴¹Ca, ⁸¹Kr and ³⁶Cl in the oceans is primarily due to the relevant thermal neutron capture reaction. Note the 50% greater in situ production of ³⁶Cl in the oceans compared to its atmospheric source.

Lal, 1999

Cosmogenic Radionuclides

Sources:

Spallation: A high energy nuclear reaction resulting in the emission of two or more fragments from the nucleus.



Cosmic Ray Shower

Cosmogenic nuclide production varies with height in the atmosphere

Most cosmic rays are “galactic” and originate from supernovas, but **our sun** is also a significant source.

89% are hydrogen (protons)
10% helium, &
1% heavier elements.


Dunai and Lifton (2014)

THE ATMOSPHERIC NUCLEAR CASCADE

P, N: high-energy (>10 MeV) secondary protons and neutrons carrying the nuclear cascade

n, p, α : secondary particles **not** carrying the nuclear cascade


π^\pm , μ^\pm , e^\pm : pions, muons, electrons, and positrons



Nuclear reactions in the atmosphere produce atmospheric cosmogenic nuclides as well as secondary particles, mostly neutrons. The latter are responsible for the cosmogenic nuclides measured in minerals. (Figure after Dunai, 2010)

Cosmogenic Radionuclides

**Cosmogenic nuclide production varies
with **latitude** in the atmosphere!**



MICHAEL OSADCIW/UNIVERSITY OF ROCHESTER

Image of solar winds deflected by Earth's magnetic field

<https://www.youtube.com/watch?v=6hD52H7rQak>




**Northern lights in County Donegal, Ireland January 24, 2012
(From a solar flare that erupted on Jan 22nd)**

Photo by Twitter user [@andrew_chester](#) via lockerz.com

^{7}Be production in the Atmosphere

Stratified



Well-mixed!





Fig. 20. A north-south section through the atmosphere showing surfaces of constant Be^7 production (nuclei/min, m^3 air at S.T.P.).

Lal and Peters (1967)

Cosmogenic nuclide production varies over seasonal to multidecadal time scales


400 Years of Sunspot Observations



"Little Ice Age"

Source variability complicates use of ^{14}C

Cosmogenic Radionuclides: Correction Factors




What are Cosmogenic Radionuclides used for?

Many studies have focused on ^{14}C (complicated by “bomb” ^{14}C and “Suess Effect”) and ^7Be given their well-defined source (atmospheric) to the surface ocean.

Studies include:

- a) Large-scale ocean circulation
 ^{14}C ($t_{1/2} = 5,700$ yr)
- b) Short time scale upper ocean vertical mixing rates and atmospheric deposition
 ^7Be ($t_{1/2} = 53$ d)
- c) Age dating, mixing, particle transport and erosion
 ^7Be ($t_{1/2} = 53$ d)
 ^{14}C ($t_{1/2} = 5,700$ y)


Cosmogenic Radionuclides: a) Ocean Circulation



stempreacademy.hawaii.edu/c-more/ocean-conveyor-belt


W. Broecker

Circulation and age of ocean waters at 3500 m determined from ^{14}C



Cosmogenic Radionuclides: c) Age Dating


Data from the Sargasso Sea



Redrawn from Druffel et al. (2016)

Cosmogenic Radionuclides: b) Upper Ocean Mixing

^{7}Be ($t_{1/2}=53\text{ d}$) as conservative tracer for upper ocean mixing: conceptual model




$$H \frac{\partial C}{\partial t} = F_a - J - \lambda C_0 H + w_H C_o - P$$

$$J = w_H C_o - K_z \left(\frac{\partial C}{\partial z} \right)_H$$

- C_o = change in ^{7}Be concentration along a mixing line (dashed)
- F_a = Atmospheric ^{7}Be input (assume constant)
- P = Particle export (assume negligible)
- H = mixed layer depth
- H_E = depth of euphotic zone
- W_H = upwelling velocity
- Dashed arrows = horizontal flow

Vertical mixing or eddy diffusivity (K_z) is derived from the vertical profile of ^{7}Be activities.



Anthropogenic Radionuclides

Bikini island 1946



Wikipedia.org

Anthropogenic Radionuclides are those radioactive nuclides that have been produced by humankind.

Major sources of Anthropogenic Radionuclides to the environment



1) Nuclear weapons



2) Nuclear fuel cycle

Why produce Anthropogenic radionuclides?

Nuclear power


It's all about using a fission chain reaction to release energy!

The main fissionable radionuclides are:

1) ^{233}U (produced by neutron irradiation of ^{232}Th)

2) ^{235}U (naturally occurring ~0.72% of U)

3) ^{239}Pu (produced by neutron irradiation ^{238}U)




Most nuclear reactors use Low Enriched Uranium (LEU) that is about 3-5% uranium-235 for ***controlled nuclear fission*** to generate heat.

Why produce Anthropogenic radionuclides?

Nuclear weapons

Reactions involved in nuclear weapons are characterized by *uncontrolled nuclear fission*.

High Enriched Uranium (HEU) is comprised of more than 20% ^{235}U and is weapons-usable, but the lower the enrichment level, the greater the amount of material required to achieve a critical mass, i.e., mass to build a nuclear bomb.



Weapons-grade HEU is typically defined as 90% HEU or above. This minimizes weapon size. Smaller and lighter nuclear weapons are much easier to deliver. Some nuclear weapons also use Pu.

Alternative uses of Anthropogenic Radionuclides

Useful for medical, scientific, and industrial applications as well as various consumer products.

Radioactive Isotope	Medical Applications
Cobalt-60	<i>Radiation therapy</i>
Iodine-131	<i>Location of brain tumors, monitoring of cardiac, liver, and thyroid activity</i>
Carbon-14	<i>Study of metabolism changes for patients with diabetes, gout, and anemia</i>
Sodium-24	<i>Studies of blood circulation</i>
mTechnecium-99	<i>Heart tissue damage, tumor detection, other medical diagnoses</i>
Industrial/Scientific Applications	
Americium-241	<i>Determination of oil wells, uniform steel and paper thickness</i>
Iridium-192	<i>Boiler and aircraft integrity</i>
Uranium-235	<i>Nuclear power</i>
Californium-252	<i>Soil moisture</i>
Consumer Products	
Americium-241	<i>Smoke Detectors</i>
Hydrogen-3 (Tritium)	<i>Watches</i>

Major sources of *known* Anthropogenic Radionuclides to the marine environment

1) Nuclear weapons - atmospheric testing

1950's-1960's most above ground tests in the Pacific Ocean, the USA and the former USSR.

$\approx 200,000 \text{ PBq}$ total; $\approx 1,000 \text{ PBq}^{137}\text{Cs}$ of which *600 PBq to oceans*.

2) Planned releases from nuclear fuel reprocessing plants

Examples: Sellafield (UK) and Cap de la Hague (France);
 $\approx 45 \text{ PBq}$ total; *$\approx 40 \text{ PBq to oceans}$ ^{137}Cs (from 1950-2010, peaking in 1975)*.

$$1 \text{ PBq} = \text{peta-Becquerel} = \text{one million billion Bq} = 10^{15} \text{ Bq}$$

IAEA (1999), IAEA (2001), and Hu et al. (2010)

Major sources of *known* Anthropogenic Radionuclides to the marine environment

3) Accidents related to nuclear power/production

- Chernobyl (April 26, 1986) \approx 5000 PBq total;
 \approx 85 PBq ^{137}Cs of which **15-20 PBq to oceans.**
- Fukushima (March 11, 2011) \approx 500 PBq total;
 \approx 20-40 PBq ^{137}Cs of which **15-20 PBq to oceans.**

4) Other main accidents with known releases in the oceans

- Satellite with nuclear power supply: SNAP-9A (1964)
 \approx **0.630 PBq** ^{238}Pu mainly in Southern Hemisphere
- Undetonated weapons in Palomares, Spain (6 Jan 1966)
 \approx **1.4 PBq** Pu isotopes
- Undetonated weapons in Thule, Greenland (21 Jan 1968)
 \approx **0.03 PBq** Pu isotopes

1 PBq = peta-Becquerel = one million billion Bq = 10^{15} Bq

IAEA (1999), IAEA (2001), and Hu et al. (2010)

Major *potential (not well constrained)* sources of Anthropogenic Radionuclides to the marine environment

1) Disposal at sea of radioactive waste (wastes from nuclear industry, medical and scientific centers)

- Known nuclear waste dumping to ocean (1946-1993)
 $\approx 85 \text{ PBq total}$

2) Accidents and losses at sea (nuclear powered submarines or vessels, undetonated nuclear weapons, nuclear powered lighthouses, sealed sources, etc.)


- $\approx 1 \text{ to } 2000 \text{ PBq}$ (e.g., Kursk submarine reactors)
- $\approx 45 \text{ PBq}$ Other categories



1 PBq = peta-Becquerel = one million billion Bq = 10^{15} Bq

IAEA (1999), IAEA (2001), and Hu et al. (2010)

Nuclear Weapons Testing 1945 - 2007



Country:	Year of first detonation:	Number of detonations:		
		atmospheric	underground	underwater
United States	1945	206	912	5
USSR	1949	223	756	3
United Kingdom	1952	21	24	
France	1960	50	160	
China	1964	22	26	
Israel?	1967?			
India	1974		6	
South Africa?	1979?	1?		
Pakistan	1998		7	
North Korea	2006		1	

not all data is official, and some locations are approximate. data source: <http://www.johnstonsarchive.net/nuclear/tests>

Each explosion is represented by a circle.
Many of these circles overlap.

Filled circles are atmospheric detonations

Hollow circles are underground or underwater tests

The size of each circle represents the yield of the blast.
The scale is not linear:

- more than 20 megatons
- 2.5–5.1 megatons
- 160–320 kilotons
- less than 15 kilotons

Map created by radicalcartography.net

Atmospheric Nuclear Weapons Testing 1945 - 1980


Atmospheric nuclear weapons testing led to temporal and spatial variability in *global* radioactive fallout by atmospheric deposition.

12% of the fallout deposited as “**close-in**” fallout near the test site

10% percent has ended up around the same latitude as the test site- “**intermediate**”

78% percent is “**global**” fallout, most of which ended up in the **same** hemisphere


Worldwide Deposition of ^{90}Sr through 1976



Data replotted from Feely et al. (1978)

Planned Releases

Discharges of major beta gamma emitters in liquid effluent from Sellafield: 1952-2015




1 TBq = tera-Becquerel = 10^{12} Bq

Adapted from Gray et al. (2005)



Source from nuclear fuel reprocessing discharges to Irish sea from Sellafield


Nuclear Power Plant Accidents (not comprehensive)



http://www.grid.unep.ch/products/4_Maps/nuclear_concernsb.jpg

Why so much attention on cesium?

Total Releases from Fukushima Dai-ichi accident




Uneven distributions of radionuclides in the ocean




Modified from Buesseler (2014)

Total cesium-137: What are the largest sources?



Total cesium-137: How does it compare to natural radioactivity?




Updated from Buesseler (2014) based on Buesseler et al. (2017)

Tracer applications of Anthropogenic Radionuclides in the ocean

1

Water/Atmospheric mixing rates
(i.e., ocean, groundwater etc.):



3

Biological uptake: ^{14}C
(see slide 16)

4

Sediment accumulation & mixing:



2

Particle scavenging & transport
rates in the water column:



5

Age dating:



Case studies

Ocean Circulation


Anthropogenic radionuclides derived from (1) atmospheric fallout (weapons tests), (2) point sources from nuclear facilities (Sellafield & La Hague) and (3) nuclear accidents (Fukushima) used to trace ocean circulation.

Particle Transport and Burial

Anthropogenic radionuclides that are particle reactive may be used to determine particle transport pathways, as well as sediment geochronologies.

1) Circulation from atmospheric fallout - weapons tests: ${}^3\text{H}$ ($t_{1/2} = 12.3$ y)


Measuring how much ${}^3\text{H}$ has penetrated into the ocean since nuclear weapons testing allows for an estimate of water mass age and circulation patterns.



1) Circulation from atmospheric fallout - weapons tests:


^{14}C ($t_{1/2} = 5730$ y)

Measuring how much ^{14}C has penetrated into the ocean since weapons testing allows for an estimate of water mass age and circulation patterns.




Povinec et al. (2009)

2) Circulation from point sources - nuclear facilities



2) Circulation from point sources - nuclear facilities

^{129}I ($t_{1/2} = 1.6 \times 10^7$ y)



Pathways (red arrows) for the transport of tracers from Sellafield and La Hague nuclear facilities through the Arctic Ocean.

Inset shows ^{129}I input functions from the nuclear fuel reprocessing plants.

● ^{129}I ($t_{1/2} = 1.6 \times 10^7$ y)

Karcher et al. (2012)


2) Circulation from point sources - nuclear facilities

$$^{129}\text{I} \quad (t_{1/2} = 1.6 \times 10^7 \text{ y})$$

^{129}I water-depth profiles in Arctic Ocean show the arrival of the large pulse produced in the 1990s from Sellafield and La Hague.

The arrival of the lower, pre-1993 signal is indicated at each station by blue ^{129}I profiles. The arrival of the higher, post-1993 signal is indicated by red profiles and provides a constraint on transit times (e.g., 10 y for transport from 60°N to St. 4 at the North Pole).

Smith et al. (2011)




2) Circulation from point sources - nuclear facilities

^{137}Cs ($t_{1/2} = 30.2 \text{ y}$)


^{137}Cs input function
(lower right inset) shows maximum values (red symbols) between 1978 and 1990.

These elevated ^{137}Cs levels appear at Sta. 4 (red lines) between 1991 and 2003 which also indicates a 10 y transit time to Sta. 4 from 60°N .

Smith et al. (2011)



3) Circulation from nuclear accidents: Fukushima, March 2011




Atmospheric Fallout to ocean:
 $\sim 15 \text{ PBq}^{137}\text{Cs}$

Direct Discharge to ocean:
 $\sim 5 \text{ PBq}^{137}\text{Cs}$




History of ^{137}Cs activity in ocean immediately off Fukushima



Kaeriyama (2016)

3) Circulation from nuclear accidents: Fukushima, March 2011



Rossi et al., 2013 & 2014

(Hover over image and click to play video)

Case studies


Ocean Circulation

Anthropogenic radionuclides derived from (1) atmospheric fallout (weapons tests), (2) point sources from nuclear facilities (Sellafield & La Hague) and (3) nuclear accidents (Fukushima) used to trace ocean circulation.

Particle Transport and Burial

Anthropogenic radionuclides that are particle reactive may be used to determine **particle transport pathways**, as well as **sediment geochronologies**.

4) Scavenging Tracers




Particle scavenging leads to decrease in atmospherically derived $^{239+240}\text{Pu}$ over time.

Subsurface peak at depth is hypothesized to be due to vertical transport followed by subsurface biological remineralization.

Lindahl et al. (2010)

4) Scavenging Tracers and Age Dating



5) Age Dating: Event Markers




FIGURE 2. ^{137}Cs activity (closed circles) and year of deposition (calculated using unsupported ^{210}Pb , open circles) plotted versus sediment depth. Samples corresponds to 0.5-cm-thick sediment slices. Years of deposition are shown with an envelope of uncertainty ($\pm 1\sigma$).

Lima et al. (2005)


Chernobyl Accident

Peak in Atmospheric Nuclear Weapons testing

Sediment core was collected from the Pettaquamscutt River Basin, Rhode Island, USA

How to find marine radioactivity data?

e.g. IAEA MARiS Database for Anthropogenic Radionuclides



<http://maris.iaea.org>

<http://maris.iaea.org>

Geotraces: <http://www.egeotrades.org/>

In summary

Cosmogenic and **anthropogenic radionuclides** are very useful tools as **tracers** to study a **large variety of processes** in the oceans at various **time scales** from days to thousands of years, such as:

- Inputs and outputs (e.g., ^{14}C , ^{137}Cs , ^3H , ^{129}I)
- Circulation (e.g., ^7Be , ^{14}C , ^{137}Cs , ^3H , ^{129}I)
- Transfer of substances (e.g., ^{14}C , $^{239+240}\text{Pu}$)
- Sedimentary record (e.g., ^7Be , ^{14}C)

Acknowledgements

This work would not have been possible without the generous contributions and thoughtful comments of Drs. Robert Anderson, Kirk Cochran, Peter Santschi and Alan Shiller. We wish to thank two anonymous reviewers who provided constructive comments that improved the presentation. Lectures would not have been possible without the outstanding assistance of graphic designer Jason Emmett.

This work was supported by the Scientific Committee on Ocean Research, SCOR Working Group 146: *Radioactivity in the Ocean, 5 Decades Later (RiO5)* and the Woods Hole Oceanographic Institution's *Center for Marine and Environmental Radioactivity*.

References

- Broecker, W.S., Peng, T-H (1982) Tracers in the sea. New York: Lamont-Doherty Geological Observatory. Eldigio Press, Columbia University. Available online at:
http://www.ideo.columbia.edu/~broecker/Home_files/TracersInTheSea_searchable.pdf
- Buesseler, K.O. (2014) Fukushima and ocean radioactivity. *Oceanography* 27(1):92–105, <http://dx.doi.org/10.5670/oceanog.2014.02>.
- Buesseler, K., Dai, M., Aoyama, M., Benitez-Nelson, C., Charmasson, S., Higley, K., Maderich, V., Masqué, P., Oughton, D., Smith, J.N. (2017) Fukushima Daiichi–Derived Radionuclides in the Ocean: Transport, Fate, and Impacts. *Annual Review of Marine Science*, 9.
- Druffel, E.R.M., Beaupré, S.R., Ziolkowski, L.A. (2016). Chapter 5: Radiocarbon in the Oceans, in Radiocarbon and Climate Change: Mechanisms, Applications and Laboratory Techniques. Editors E.A.G. Schuur, S.E. Trumbore and E.R.M. Druffel, Springer International Publishing, New York/Heidelberg, 139-166.
- Dunai, T.J., Lifton, N.A. (2014) The Nuts and Bolts of Cosmogenic Nuclide Production. *Elements Magazine*, 10(5):347-350, 1811-5209/14/0010-347\$2.50 DOI: 10.2113/gselements.10.5.347.
- Freely, H.W., Volchok, H.L., Hardy, E.P., Toonkel, L.E. (1978) Worldwide deposition of ⁹⁰Sr through 1976. *Environment Science and Technology*, 12: 808–809.
- Gray, J., Jones, S.R., Smith, A.D. (1995) Discharges to the environment from the Sellafield site, 1951-1992. *Journal of Radiological Protection*, 15, 99–131.
- Hancock, G., Tims, SJ, Fifield, LF, Webster, IT (2014) The release and persistence of radioactive anthropogenic nuclides. Geological Soc., London Special Publications. 395: 265-281 doi: 10.1144/SP395.15
- Haskell, W.Z., Kadko, D., Hammond, D.E., Knapp, A.N., Prokopenko, M.G., Berelson, W.M., Capone, D.G. (2015). Upwelling velocity and eddy diffusivity from ⁷Be measurements used to compare vertical nutrient flux to export POC flux in the Eastern Tropical South Pacific. *Marine Chemistry*, 168: 140-150.
- Hu, Q-H, Weng, J-Q, Wang, J-S (2010) Sources of anthropogenic radionuclides in the environment: a review. *Journal of Environmental Radioactivity* 101: 426-437.
- International Atomic Energy Agency (1999) Inventory of Radioactive Waste Disposals at Sea, IAEA TECDOC No. 1105.
- International Atomic Energy Agency (2001) Inventory of Accidents and Losses at Sea Involving Radioactive Material, IAEA TECDOC No. 1242.

References

- Kadko, D., Johns, W. (2011) Inferring upwelling rates in the equatorial Atlantic using ^{7}Be measurements in the upper ocean. Deep Sea Research Part 1: Oceanographic Research Papers, 58: 647-657. doi:10.1016/j.dsr.2011.03.004
- Kaeriyama, K. (2016) Oceanic dispersion of Fukushima-derived radioactive cesium: a review. Fisheries Oceanography, 1-15; doi:10.1111/fog.12177
- Karcher, M., Smith, J.N., Kauker, F., Gerdes, R., Smethie Jr., W.M. (2012) Recent changes in Arctic Ocean circulation revealed by iodine-129 observations and modeling, Journal of Geophysical Research, 117; C08007, doi: 10.1029/2011JC007513.
- Lal, D. (1999) An overview of five decades of studies of cosmic ray produced nuclides in oceans. Sci. of the Total Environ., 238: 3-13.
- Lal, D., Peters, B. (1967) Cosmic ray produced radioactivity on the earth. In: S. Flugg (Editor), Handbuch der Physik. Springer, Berlin, pp. 551-612.
- Lima, A.L., Hubeny, J.B., Reddy, C.M., King, J.W., Hughen, K.A., Eglinton, T.I. (2005). High-resolution historical records from Pettaquamscutt River basin sediments: 1. ^{210}Pb and varve chronologies validate record of ^{137}Cs released by the Chernobyl accident. Geochimica et Cosmochimica Acta, 69(7); 1803-1812.
- Lindahl, P., Lee, S-H, Worsfold, P., Keith-Roach, M. (2010) Plutonium isotopes as tracers for ocean processes: A review. Marine Environmental Research, 69: 73-84.
- Matsumoto, K., Key, R. (2004) Natural radiocarbon distribution in the deep ocean, in Global Environmental Change in the Ocean and on Land, edited by M. Shiomi, and H. Kawahata, TERRAPUB, Tokyo, 45–58.
- Povinec, P.P., Litherland, A.E., von Reden, K.F. (2009) Developments in Radiocarbon technologies from the Libby Counter to compounds specific AMS Analyses, 51(1):45–78.
- Reimer, P.J., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Bronk, C., Buck, C.E., Cheng, H., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Haflidason, H., Hajdas, I., Hatté, C., Heaton, T.J., Hoffmann, D.L., Hogg, A.G., Hughen, K.A., Kaiser, K.F., Kromer, B., Manning, S.W., Niu, M., Reimer, R.W., Richards, D.A., Scott, E.M., Southon, J.R., Staff, R.A., Turney, C.S.M., van der Plicht, J. (2013) IntCal13 and Marine13 radiocarbon age calibration curves 0–50,000 years cal BP. Radiocarbon, 55(4):1869–1887.
- Richards, Z.T., Beger, M., Pinca, S., Carden, Wallace, C. (2008) Bikini Atoll coral biodiversity resilience five decades after nuclear testing. Marine Pollution Bulletin, 56 (3): 503–515. doi:10.1016/j.marpolbul.2007.11.018.

References

- Rossi, V., Van Sebille, E., Gupta, A.S., Garçon, V., England, M.H. (2013) Multi-decadal projections of surface and interior pathways of the Fukushima Cesium-137 radioactive plume. Deep Sea Research Part I: Oceanographic Research Papers 80: 37-46.
- Rossi, V., Sebille, E.V., Gupta, A.S., Garçon, V., England, M.H. (2014) Corrigendum to “Multi-decadal projections of surface and interior pathways of the Fukushima Cesium-137 radioactive plume” [Deep-Sea Res. I 80 (2013) 37–46], Deep-Sea Research, Part I, 93: 162–164.
- Rutgers van der Loeff, M.M. (2014) Uranium-Thorium Decay Series in the Oceans: Overview. In Elias S (ed.) Earth Systems and Environmental Sciences, (Reference Module in Earth Systems and Environmental Sciences). Amsterdam: Elsevier 1-16.
- Smith, J.N., McLaughlin, F.A., Smethie Jr., W.M., Moran, S.B., Lepore, K. (2011), Iodine-129, 137Cs, and CFC-11 tracer transit time distributions in the Arctic Ocean, Journal of Geophysical Research, 116: C04024, doi: 10.1029/2010JC006471.

Copyright

**Copyright ©2018 by The Association for the Sciences
of Limnology and Oceanography (ASLO)**

Published by John Wiley & Sons, Inc.

**Claudia Benitez-Nelson, Ken Buesseler, Minhan Dai,
Michio Aoyama, Núria Casacuberta, Sabine Charmasson,
José Marcus Godoy, Andy Johnson, Vladimir Maderich,
Pere Masqué, Willard Moore, Paul J. Morris, John N. Smith**

***Radioactivity in the Marine Environment:
Cosmogenic and Anthropogenic Radionuclides***

doi:10.1002/loe2.10009

e-Lecture received June 2016; accepted October 2018

**For more information, please visit the ASLO e-Lectures page at
[http://onlinelibrary.wiley.com/journal/10.1002/\(ISSN\)2164-0254](http://onlinelibrary.wiley.com/journal/10.1002/(ISSN)2164-0254)**