

ATMOSPHERIC TRANSPORT MODELLING OF RADON-222 AT EUROPEAN SCALE: DESCRIPTION AND VALIDATION AGAINST ICOS OBSERVATIONS.

Arnaud Quéré¹ , Tanina Hached¹ , Denis Quélo¹ , Michel Ramonet² , Camille Yver Kwok² ,
and Ute Karstens³

¹ IRSN, Fontenay-aux-Roses cedex, France (arnaud.querel@irsn.fr)

² LSCE, Gif-sur-Yvette, France (michel.ramonet@lsce.ipsl.fr)

³ Lund University, Lund, Sweden (ute.karstens@nateko.lu.se)

Context


- LdX is an atmospheric transport model (**ATM**) used for assessing the consequences of nuclear accidents
 - Validated against the Fukushima accident (<https://doi.org/10.1016/j.jenvrad.2021.106712>)
 - And on the radon progeny wet deposition (<https://doi.org/10.5194/adgeo-57-109-2022>)
- Compared with the observed radon air concentrations, the radon modelling framework can be used to:
 - Validate the modelled radon-222 air concentrations
 - Help to screen the radon exhalation flux maps
- Study partially funded by the RADONORM project, under EURATOM Horizon 2020
 - <https://www.radonorm.eu/>
- In collaboration with the traceRadon project, co-financed by the Participating States and from the European Union's Horizon 2020
 - <http://traceradon-empir.eu/>



Simulation framework

Input

- Radon exhalation flux



- Physical properties (decay chain, ...)

- Meteorological data



ICOS Input

- ICOS network Rn-222 air concentration

trace
Radon

Processing

Atmospheric transport model (Ldx)




- Eulerian long-range transport model
- Mainly used in nuclear emergency responses


Processing

Model/measure comparison

- Rn-222 air concentration time series



- Air concentration maps (Bq/m3)



- Data for further analyses

ICOS_score_global.csv
ICOS_sequential_D.csv
ICOS_sequential_M.csv
ICOS_sequential_W.csv
Score_global.csv
Score_peak_success.csv
Score_peak_success_permiss...
Summary_all_peaks.csv

ICOS network

ICOS
Integrated Carbon Observation System




<https://www.icos-cp.eu/>

Heiskanen et al. (2021), <https://doi.org/10.1175/BAMS-D-19-0364.1>

9 stations available for 2021

ID	Station name, Country	Heigh agl (m)	Ground altitude asl (m)	Mean obs value in 2021 (Bq/m³)
MHD	Mace Head, Ireland	24	5	0,3
SAC2	Saclay, France	2	160	1.5
SAC100	Saclay, France	100	160	2.4
OPE	OPE, France	120	390	3.1
TRN	Trainou, France	180	131	2.6
PUY	Puy-de-Dôme, France	10	1465	0.3
PDM	Pic-du-Midi, France	10	2877	2.4
INTE	Barcelona, Spain	5	65	2.3
FKL	Finokalia, Greece	10	250	0.5



Exhalation rate with GLDAS soil moisture 20 Dec 2020, 00:00

Radon exhalation flux maps

Exhalation flux based on Karstens et al.
(2015)
<https://doi.org/10.5194/acp-15-12845-2015>


Inputs :

- uranium content,
- soil texture,
- Porosity
- soil moisture

Updated until 2022 with new ERA5 and
GLDAS Noah soil moistures

Karstens, U. and Levin, I. (2022). traceRadon
monthly radon flux map for Europe 2006-
2022 (based on GLDAS-Noah v2.1 soil
moisture)




https://hdl.handle.net/11676/ge5vMeklvG_Qz43rzS2wx0-



Radon exhalation

3 different exhalation maps are compared:

Karstens 2022 ERA5	Karstens 2022 GLDAS	Karstens 2015 GLDAS
Calculated in 2022 with the monthly mean soil moisture estimates ERA-Interim/Land model.	Calculated in 2022 with the monthly mean soil moisture estimates from the GLDAS.	Calculated in 2015 with the monthly mean soil moisture estimates from the GLDAS Noah. 2021 exhalation is extrapolated from the monthly mean during the period.



Visual results : time series


Rn-222 air conc. at SAC_100 (Saclay, France)
resampled on hourly frequency

■ Radon exhalation map :
Karstens 2022 GLDAS

■ 01 jan 2021-> 31 dec 2021

■ Hourly frequency


Good results on base
line and frequency



Visual results : time series

- Radon exhalation map : Karstens 2022 GLDAS
- 01 jan 2021-> 31 dec 2021
- Week frequency
- PUY and PDM badly modelled, for all exhalation map.

ATM not designed to deal with the mountainous stations




Statistics, focus on the wind speed

Radon exhalation map : Karstens
2022 GLDAS

01 jan 2021-> 31 dec 2021



Good correlation between sim and obs.

For both sim and obs, a low wind speed is correlated with a high Rn-222 concentration.




Statistics, focus on the wind speed

- Radon exhalation map : Karstens 2022 GLDAS
- 01 jan 2021-> 31 dec 2021





Statistics




Correlation
Worst 55 Better 80 100

SAC 0 m agl	SAC 100 m agl	OPE 120 m agl	T 180
83	88	74	
61		55	
81	84	70	




Large variations between each stations.

Statistics



Normalised RMSE point of view

- A soil moisture estimated from ERA5 seems to lead to a better exhalation map for stations lower than 25 m agl
 - Issue with mean bias in GLDAS or ERA5 exhalation ?
 - Issue with the vertical transport of the ATM ?
 - Saclay local features ?





Statistics

- Radon exhalation map : Karstens 2022 GLDAS
- 01 jan 2021-> 31 dec 2021

Oddly, no correlation between obs and exhalation !


kz simu (% heure >1
wind module
wind angle
pressure
temperature



Statistics

- Radon exhalation map : Karstens 2022 GLDAS
- 01 jan 2021-> 31 dec 2021

**Oddly, no correlation between obs and exhalation !
(found on 8/9 of the station)**



Conclusion

- Keys to evaluate the radon flux maps
 - The ERA5 soil humidity leads to worse results
 - Downside: there are few stations, and few comparable stations
- Wind is the key parameter to the radon air concentration

Outlook



- Extend the study to the AGES stations
 - RADONORM project (WP2.2)
- Odd results about exhalation/observed concentrations correlation.
- Feed back on exhalation
- More stations, please !

Thank you for your attention !